



Cambridge International Examinations
Cambridge Ordinary Level

PHYSICS

5054/22

Paper 2 Theory

May/June 2016

MARK SCHEME

Maximum Mark: 75

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2016 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

© IGCSE is the registered trademark of Cambridge International Examinations.

This document consists of **6** printed pages.

© UCLES 2016



Page 2	Mark Scheme	Syllabus	Paper
	Cambridge O Level – May/June 2016	5054	22

- 1 (a) mark at a time between 4.0 and 7.5 seconds B1
- (b) (a =) $(v - u)/t$ numerical or algebraic C1
or (a =) gradient of graph stated
2.5 m/s² A1
- (c) forward force and backward force clear B1
equal forces (in horizontal direction) B1
or no resultant force
or forces cancel/balance/in equilibrium
- 2 (a) (PE =) mgh **or** Fd **or** 5×3.5 C1
17.5 J **or** 17 J **or** 18 J A1
- (b) (i) (efficiency = useful) energy output/energy input B1
in any form but all three quantities must be mentioned
if efficiency is not the subject of the equation
- (ii) 17.5/0.65 **or** 17.5/65 C1
or $0.65/65 = \text{(a)}/\text{energy input}$
26.9 J **or** 27 J A1
- (c) due to friction (in bearings of motor) B1
or due to (electrical) resistance (in motor)
or air resistance acts
or thermal energy/heat produced/lost (in resistance of motor/due to friction)
- 3 (a) (i) C M1
- (ii) **data** quoted to prove stretches more at end A1
or extensions/changes in length increase/are not the same (at higher loads)
- (iii) 4.5 cm B1
- (b) (tie rock to spring A) B1
- find weight/force/newtons using length or extension **and** graph **or** match readings (in table)
 - find known weight/mass/force/N that gives same extension of spring
 - use of proportionality with length or extension
 - extension (in cm)/1.6
- (mass =) weight/g B1
or weight/gravitational field (strength)

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge O Level – May/June 2016	5054	22

- 4 (a) steam B1
or (water) vapour
or water in gaseous form
- (b) (E =) mL numerical or algebraic C1
or 52 000–6000 or 46 000 (J) seen
- (52 000–6000)/20 or 46 000/20 C1
2300 J/g or 2.3×10^6 J/kg A1
- (c) fast moving/energetic molecules escape/evaporate/break bonds/become gas B1
leaving slow(er) molecules/less energetic molecules B1
or reducing **average** (kinetic) energy (of molecules or liquid)
- 5 (a) (i) long-sight or far-sight or hypermetropia B1
- (ii) rays do not come together (on back of eye) B1
or rays do not converge (on retina)
or it/the image is not formed on retina/back of eye
or it/the image is formed behind retina/back of eye
- (b) (i) lens between rays and eyeball **and** a converging lens shown B1
- (ii) converging or convex B1
- 6 (a) (i) red B1
- (ii) blue B1
- (b) ANY 2 from (the use must agree with the type) B4
- Microwaves B1
use – satellite television, telephone, mobile/cell phones; B1
cooking, heating in a microwave oven, television
remote, radar, communication
- X(-rays) B1
use – hospital use in medical imaging or security imaging, killing cancerous cells, B1
and engineering applications such as detecting cracks in metal, crystallography
- gamma (rays) B1
use – medical treatment in killing cancerous cells, and engineering applications B1
such as detecting cracks in metal, sterilisation, tracer applications, radiotherapy

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge O Level – May/June 2016	5054	22

- 7 (a) any insulator, e.g. perspex, plastic, nylon, rubber B1
- (b) top of P shows a net negative charge with some negative charges under rod B1
bottom of P has equal number of positive charges B1
- (c) (i) clear net negative charge on P B1
and (net) negative charges above or at middle line
- (ii) 1 negative (charges)/electrons flow to earth B1
or (P) becomes neutral
- 2 charges spread over P B1
- 8 (a) current/a.c (in primary coil) creates magnetic field B1
or current/a.c magnetises iron B1
changing magnetic field (in secondary)
- (b) it/secondary has less turns (than primary) B1
or primary has more turns (than secondary)
or (some) flux escapes
- (c) (steel is) a permanent magnet B1
or weaker fields produced
or (steel) difficult to magnetise/demagnetise
or (steel) is a hard magnetic material
- (d) passes current/charge in one direction B1
or has high resistance/is an insulator when current in
one direction/reverse biased
- 9 (a) (amount of) energy/work (dissipated by source) M1
by unit charge (around a circuit) A1
- (b) (i) 1 they are the same B1
or $I_B = I_1 = I_2$
- 2 $E = V_1 + V_2$ B1
- (ii) (I=) V/R in any form algebraic or numerical C1
0.25 A A1
- (iii) 4.5 V B1
- (iv) (P=) VI **or** (P=) I^2R **or** (P=) V^2/R C1
in any form algebraic or numerical
1.1(25) W A1

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge O Level – May/June 2016	5054	22

- (c) current is (directly) proportional to voltage B1
or voltage/current is a constant
law holds for constant physical conditions/ B1
constant temperature/constant pressure/for metals
- (d) (i) (directly) proportional B1
or $(R) \propto 1$
- (ii) inversely proportional B1
or $(R) \propto 1/A$
- (e) 1st band orange B1
2nd and 3rd bands both black B1
- 10 (a) (i) B – anode B1
D – filament or heater B1
E and F–Y plates or X plates in either order B1
- (ii) 1 attract electrons or gives electrons speed/K.E. B1
- 2 heats up cathode B1
or gives electrons energy to escape (metal/cathode)
or causes/allows thermionic emission
- (iii) kinetic energy to light B1
or electrical energy to light
- (iv) voltage/charge is applied to the X-plates/vertical plates B1
or turn on time base
(steadily) increasing voltage/charge applied to plate(s) B1
or saw tooth voltage applied
or electrons attracted/repelled by plate(s) or by the electric field between them
- (b) (i) 1 1(.0)V B1
- 2 one wave 1.3–1.4 squares or 3 waves in 4 squares C1
2.6–2.8 ms A1
- 3 $(f =) 1/T$ numerical or algebraic C1
345–400 Hz A1
- (ii) smaller amplitude shown B1
larger period shown B1

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge O Level – May/June 2016	5054	22

- 11 (a) (nucleus/nuclide/atom) with same number of protons B1
- (b) (i) 2 B1
- (ii) neutron B1
- (iii) 2 B1
- (iv) 4 B1
- (c) nuclei repel B1
or like/positive charges repel
(needs) high kinetic energy/speed (to overcome repulsion) B1
- (d) ANY 3 lines from B3
(dust/gas) collapses/comes together/clusters/condenses
gravitational attraction or gravity mentioned B1
temperature rises or KE (dust/gas) increases B1
(nuclear) fusion occurs B1
equilibrium established as radiation pressure/outward
force balances inward force B1
- (e) (i) time for a quantity to halve C1
time for (radio)activity/count rate/number of atoms/number of nuclei to halve A1
- (ii) any relevant halving seen, e.g. 16 000/2 C1
1000 A1