MARK SCHEME for the October/November 2013 series

9709 MATHEMATICS

9709/21

Paper 2 (Pure Mathematics), maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.





Page 2	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – October/November 2013	9709	21

Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Mark for a correct result or statement independent of method marks.
- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol s^h implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0. B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking *g* equal to 9.8 or 9.81 instead of 10.



Page 3	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – October/November 2013	9709	21

The following abbreviations may be used in a mark scheme or used on the scripts:

- AEF Any Equivalent Form (of answer is equally acceptable)
- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
- CWO Correct Working Only often written by a 'fortuitous' answer
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)
- SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through √" marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.



	Page 4			Mark Schen		Syllabus	Paper	r
			GCE A	S LEVEL – October	/November 2013	9709	21	
1	<u>Either</u>	State or imply non-modular inequality $(x+1)^2 < (3x+5)^2$, or corresponding equation or pair of linear equations Make reasonable solution attempt at a 3-term quadratic, or solve						
		two]	inear equatio		term quadratic, or solve		M1 A1	
				yer $x < -2$ or $x > -\frac{3}{2}$			A1	
	<u>Or</u>	or fro Obta	om a graphica in the other c	l value, e.g. $x = -2$, by al method or by inspect ritical value similarly for $x < -2$ or $x > -\frac{3}{2}$	solving a linear equatio tion	n (or inequality)	B1 B2 B1	[4]
2	(i)	Com	plete the argu	$x^4 + 2x - 9$ at $x = 1.5$ at ment correctly with ap f(1.6) = 0.7536			M1 A1	[2]
	(ii) Rearrange $x^4 + 2x - 9 = 0$ to given equation or vice versa				B1	[1]		
	(iii)	Obta	in final answ	ormula correctly at lea er 1.56 erations to justify its ac			M1 A1 B1	[3]
			= 1.5 1.5874 1.5424 1.5653 1.5536	$ x_0 = 1.55 \\ 1.5614 \\ 1.5556 $	$ x_0 = 1.6 \\ 1.5362 \\ 1.5685 \\ 1.5520 \\ 1.5604 $			

or show there is a sign change in the interval (1.555, 1.565)

1 5595

1.5565

3	Obtain derivative $e^{2x} - 5e^x + 4$	B1	
	Equate derivative to zero and carry out recognisable solution method for a quadratic in e^x	M1	
	Obtain $e^x = 1$ or $e^x = 4$	A1	
	Obtain $x = 0$ and $x = \ln 4$	A1	
	Use an appropriate method for determining nature of at least one stationary point	M1	
	$\left(\frac{d^2 y}{dx^2} = 2e^{2x} - 5e^x, \text{ when } x = 0, \frac{d^2 y}{dx^2} = -(3), x = \ln 4, \frac{d^2 y}{dx^2} = +(12)\right)$		
	Conclude maximum at $x = 0$ and minimum at $x = \ln 4$ (no errors seen)	A1	[6]

1.5561

4	(i)	Substitute $x = 3$ and equate to 14 $(9a + 3b + 35 = 14)$	M1	
		Substitute $x = -2$ and equate to 24 $(4a - 2b = 24)$	M1	
		Obtain a correct equation in any form	A1	
		Solve a relevant pair of equations for a or for b	M1	
		Obtain $a = 1$ and $b = -10$	A1	[5]



	Page 5		5 Mark Scheme Syllabus	B Paper	er	
			GCE AS LEVEL – October/November 2013 9709	21		
	(ii) Attempt division by $x^2 + 2x - 8$ and reach a partial quotient of $x - k$ Obtain quotient $x - 1$ with no errors seen (can be done by observation) Correct solution method for quadratic e.g. factorisation					
			All solutions $x = 1$, $x = 2$ and $x = -4$ given and no others CWO	M1 A1	[4]	
5		(i)	State $\frac{dx}{d\theta} = -2\sin 2\theta + \sin \theta$ or $\frac{dy}{d\theta} = 8\sin \theta \cos \theta$	B1		
			Use $\frac{dy}{dx} = \frac{dy}{d\theta} \div \frac{dx}{d\theta}$	M1		
			Use $\sin 2\theta = 2\sin\theta\cos\theta$	M1		
			Obtain given answer correctly	A1	[4]	
		(ii)	Equate derivative to -4 and solve for $\cos \theta$	M1		
			Obtain $\cos \theta = \frac{1}{2}$ Obtain $x = -1$	A1 A1		
			Obtain $y = 3$	A1 A1	[4]	
			ooum y 5	711	[,]	
6	(a)	(i)	Attempt to divide by e^{2x} and attempt to integrate 2 terms	M1		
			Integrate a term of form ke^{-2x} correctly Fully correct integral $x - 3e^{-2x} (+c)$	A1√ [≜] A1	[3]	
		(ii)	State correct expression $\frac{1}{2}\cos 2x + \frac{1}{2}$ or equivalent	B1		
		(11)	Integrate an expression of the form $a + b \cos 2x$, where $ab \neq 0$, correctly	M1		
			State correct integral $\frac{3\sin 2x}{4} + \frac{3x}{2}(+c)$	A1	[3]	
	(b)	Stat	te or imply correct ordinates 5.46143, 4.78941, 4.32808	B1		
	(-)		e correct formula, or equivalent, correctly with $h = 0.5$ and three ordinates	M1		
		Obt	tain answer 4.84 with no errors seen	A1	[3]	
7		(i)	State $R = \sqrt{10}$	B1		
			Use trig formula to find α	M1		
			Obtain $\alpha = 18.43^{\circ}$ with no errors seen	A1	[3]	
		(ii)	Carry out evaluation of $\cos^{-1}\left(\frac{2}{R}\right) \approx 50.77^{\circ}$	M1		
			Carry out correct method for one correct answer	M1		
			Obtain one correct answer e.g. 34.6°	Al		
			Carry out correct method for a further answer	M1		
			Obtain remaining 3 answers 163.8°, 214.6°, 343.8° and no others in the range	e Al	[5]	

