

Cambridge
International
AS & A Level

Cambridge International Examinations
 Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

9701/35

Paper 3 Advanced Practical Skills 1

May/June 2016

2 hours

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
 Give details of the practical session and laboratory where appropriate, in the boxes provided.
 Write in dark blue or black pen.
 You may use an HB pencil for any diagrams or graphs.
 Do not use staples, paper clips, glue or correction fluid.
 DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.
 Electronic calculators may be used.
 You may lose marks if you do not show your working or if you do not use appropriate units.
 Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 14 and 15.
 A copy of the Periodic Table is printed on page 16.

At the end of the examination, fasten all your work securely together.
 The number of marks is given in brackets [] at the end of each question or part question.

Session
Laboratory

For Examiner's Use	
1	
2	
3	
Total	

This document consists of **13** printed pages and **3** blank pages.



- 1 In this experiment you will determine the concentration of a solution of sulfuric acid by titration.

FA 1 is sulfuric acid, H_2SO_4 .

FA 2 is aqueous sodium hydroxide, containing 4.20 g NaOH dissolved in 1.00 dm^3 of water.
 thymolphthalein indicator

(a) Method

Dilution of FA 1

- Pipette **10.0 cm^3** of **FA 1** into the 250 cm^3 volumetric flask.
- Make the solution up to the mark using distilled water.
- Shake the flask thoroughly.
- This diluted solution of sulfuric acid is **FA 3**. Label the flask **FA 3**.

Titration

- Fill the burette with **FA 2**.
- Pipette **25.0 cm^3** of **FA 3** into a conical flask.
- Add a few drops of thymolphthalein indicator.
- Perform a rough titration and record your burette readings in the space below. The end point is reached when the solution turns a permanent pale blue colour.

The rough titre is cm^3 .

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make sure any recorded results show the precision of your practical work.
- Record in a suitable form below all of your burette readings and the volume of **FA 2** added in each accurate titration.

Keep solution FA 1 for use in Questions 2 and 3.

I	
II	
III	
IV	
V	
VI	
VII	

[7]

- (b) From your accurate titration results, obtain a suitable value for the volume of **FA 2** to be used in your calculations.
 Show clearly how you obtained this value.

25.0 cm³ of **FA 3** required cm³ of **FA 2**. [1]

(c) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

- (i) Calculate the number of moles of sodium hydroxide present in the volume of **FA 2** calculated in (b).
 Use the data in the Periodic Table on page 16.

moles of NaOH = mol

- (ii) Complete the equation for the reaction of sulfuric acid with sodium hydroxide. State symbols are required.

..... + → Na₂SO₄(aq) +

- (iii) Use your answers to (i) and (ii) to calculate the number of moles of sulfuric acid used in each titration.

moles of H₂SO₄ = mol

- (iv) Calculate the concentration, in mol dm⁻³, of sulfuric acid in **FA 3**.

concentration of H₂SO₄ in **FA 3** = mol dm⁻³

- (v) Calculate the concentration, in mol dm⁻³, of sulfuric acid in **FA 1**.

concentration of H₂SO₄ in **FA 1** = mol dm⁻³
 [5]

[Total: 13]

- 2 In this experiment you will determine the enthalpy change, ΔH , for the decomposition of magnesium carbonate to magnesium oxide.



In order to do this, you will determine the enthalpy changes for the reactions of magnesium carbonate and magnesium oxide with sulfuric acid. Excess of the two magnesium compounds will be used in each experiment.

Then you will use Hess' Law to calculate the enthalpy change for the reaction above.

FA 1 is sulfuric acid, H_2SO_4 .

FA 4 is magnesium carbonate, MgCO_3 .

FA 5 is magnesium oxide, MgO .

- (a) Determination of the enthalpy change for the reaction of magnesium carbonate, **FA 4**, with sulfuric acid, **FA 1**

(i) **Method**

- Support the plastic cup inside the 250 cm^3 beaker.
- Use a measuring cylinder to transfer 25 cm^3 of **FA 1** into the plastic cup.
- Measure and record the initial temperature of the **FA 1** in the space below.
- Add all the **FA 4** from the container to the **FA 1** in the plastic cup.
- Stir constantly until the maximum temperature is reached.
- Measure and record the maximum temperature of the contents of the cup.
- Rinse out the plastic cup and shake to dry for use in (b).
- Calculate and record the temperature rise.

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

- energy produced = J

- (If you were unable to calculate the concentration of sulfuric acid in **FA 1**, assume that it is 1.27 mol dm^{-3} . This is not the true value.)

moles of H_2SO_4 = mol

- (iv) Calculate the enthalpy change, in kJ mol^{-1} , for the reaction below.



enthalpy change = kJ mol⁻¹
(sign) (value)

[6]

(i) Method


- Use the measuring cylinder to transfer approximately 40 cm³ of **FA 1** into the **100 cm³** beaker.
- Place the beaker on a tripod and gauze.
- Heat **FA 1** in the beaker until the temperature is between 40 °C and 50 °C.
- Support the plastic cup in the **250 cm³** beaker.
- Use the measuring cylinder to transfer 25 cm³ of hot **FA 1** into the plastic cup. **CARE.**
- Measure and record, in the space below, the initial temperature of **FA 1** in the plastic cup.
- Immediately, add all the **FA 5** from the container to the **FA 1** in the plastic cup.
- Stir constantly until the maximum temperature is reached.
- Measure and record the maximum temperature.
- Calculate and record the temperature rise.

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(ii) Calculate the energy produced during this reaction.
[Assume that 4.2 J are needed to raise the temperature of 1.0 cm³ of solution by 1.0 °C.]

energy produced = J

(iii) Use your answer to (a)(iii) to calculate the enthalpy change, in kJ mol^{-1} , for the reaction below.



enthalpy change = kJ mol⁻¹
(sign) (value)

[4]

- (If you were unable to calculate the enthalpy changes, assume that the value of the enthalpy change in **(a)(iv)** is $-58.7 \text{ kJ mol}^{-1}$ and the value in **(b)(iii)** is $-140.3 \text{ kJ mol}^{-1}$. Note: these are not the correct values.)



enthalpy change = kJ mol⁻¹
(sign) (value)

[2]

- (d) (i)** Calculate the maximum percentage error in the temperature **rise** in **(b)(i)**.

percentage error = %

- (ii) The magnesium oxide, **FA 5**, was weighed with a balance measuring to one decimal place. A student suggested that the accuracy of the experiment in **(b)(i)** would be improved by weighing **FA 5** using a balance measuring to two decimal places. State and explain whether or not the student is correct.

.....

.....

.....

[2

[Total: 14]

3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations.**

You should indicate clearly at what stage in a test a change occurs.

No additional tests for ions present should be attempted.

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

- (a) (i) **FA 6** is a salt containing one cation and one anion from those listed on pages 14 and 15. Transfer a **small** spatula measure of **FA 6** into a hard-glass test-tube. Heat gently at first, then heat strongly until no further change occurs.

Record **all** your observations below.

.....

.....

.....

.....

.....

.....

.....

- (ii) Dissolve the remainder of **FA 6** in an approximate depth of 5 cm of distilled water in a boiling tube for use in the following tests. Record your observations in the table below.

<i>test</i>	<i>observations</i>
To a 1 cm depth of the solution of FA 6 in a test-tube, add an equal volume of FA 1 , aqueous sulfuric acid.	
To a 1 cm depth of the solution of FA 6 in a test-tube, add aqueous ammonia.	
To a 1 cm depth of the solution of FA 6 in a boiling tube, add aqueous sodium hydroxide, then	
heat the mixture, gently and carefully, then	
place the boiling tube in a rack and add aluminium foil.	

- (iii) Give the chemical formula of **FA 6**.

.....

Give the ionic equation for the reaction of **FA 6** with cold sodium hydroxide. Include state symbols.

.....

[8]

- (b) (i) **FA 7** is a solution containing one cation and one anion from the list on pages 14 and 15.

Carry out the following tests and record your observations in the table below.

<i>test</i>	<i>observations</i>
To a 1 cm depth of FA 7 in a test-tube, add aqueous sodium hydroxide.	
To a 1 cm depth of FA 7 in a test-tube, add aqueous ammonia.	
To a 1 cm depth of FA 7 in a test-tube, add a few drops of acidified potassium manganate(VII), followed by a few drops of aqueous starch.	

- (ii) Identify **FA 7**.

FA 7 is

- (iii) Carry out **one** further test of your choice to confirm the identity of the **anion** in **FA 7**.

reagent(s) used

observation(s)

.....

[5]

[Total: 13]

Qualitative Analysis Notes

Key: [ppt. = precipitate]

1 Reactions of aqueous cations

ion	reaction with	
	NaOH(aq)	NH ₃ (aq)
aluminium, Al ³⁺ (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH ₄ ⁺ (aq)	no ppt. ammonia produced on heating	—
barium, Ba ²⁺ (aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.
calcium, Ca ²⁺ (aq)	white ppt. with high [Ca ²⁺ (aq)]	no ppt.
chromium(III), Cr ³⁺ (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg ²⁺ (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn ²⁺ (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess

2 Reactions of anions

<i>ion</i>	<i>reaction</i>
carbonate, CO_3^{2-}	CO_2 liberated by dilute acids
chloride, $\text{Cl}^-(\text{aq})$	gives white ppt. with $\text{Ag}^+(\text{aq})$ (soluble in $\text{NH}_3(\text{aq})$)
bromide, $\text{Br}^-(\text{aq})$	gives cream ppt. with $\text{Ag}^+(\text{aq})$ (partially soluble in $\text{NH}_3(\text{aq})$)
iodide, $\text{I}^-(\text{aq})$	gives yellow ppt. with $\text{Ag}^+(\text{aq})$ (insoluble in $\text{NH}_3(\text{aq})$)
nitrate, $\text{NO}_3^-(\text{aq})$	NH_3 liberated on heating with $\text{OH}^-(\text{aq})$ and Al foil
nitrite, $\text{NO}_2^-(\text{aq})$	NH_3 liberated on heating with $\text{OH}^-(\text{aq})$ and Al foil; NO liberated by dilute acids (colourless $\text{NO} \rightarrow$ (pale) brown NO_2 in air)
sulfate, $\text{SO}_4^{2-}(\text{aq})$	gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (insoluble in excess dilute strong acids)
sulfite, $\text{SO}_3^{2-}(\text{aq})$	gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (soluble in excess dilute strong acids)

3 Tests for gases

<i>gas</i>	<i>test and test result</i>
ammonia, NH_3	turns damp red litmus paper blue
carbon dioxide, CO_2	gives a white ppt. with limewater (ppt. dissolves with excess CO_2)
chlorine, Cl_2	bleaches damp litmus paper
hydrogen, H_2	"pops" with a lighted splint
oxygen, O_2	relights a glowing splint

Group																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																	
1	2	Key										13	14	15	16	17	18																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																
		atomic number atomic symbol name relative atomic mass																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															
3	4																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																
Li lithium 6.9	Be beryllium 9.0	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57–71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89–103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	53

57	La	58	Ce	59	Pr	60	Nd	61	Pm	62	Sm	63	Eu	64	Gd	65	Tb	66	Dy	67	Ho	68	Er	69	Tm	70	Yb	71	Lu
lanthanum	138.9	cerium	140.1	praseodymium	140.9	neodymium	144.4	promethium	—	samarium	150.4	europtium	152.0	gadolinium	157.3	terbium	158.9	dysprosium	162.5	holmium	164.9	erbium	167.3	thulium	168.9	ytterbium	173.1	lutetium	175.0
89	Ac	90	Th	91	Pa	92	U	93	Np	94	Pu	95	Am	96	Cm	97	Bk	98	Cf	99	Es	100	Fm	101	Md	102	No	103	Lr
actinium	227.0	thorium	232.0	protactinium	231.0	uranium	238.0	neptunium		plutonium		americium		curium		berkelium		californium		einsteinium		fermium		meitnerium		nobelium		lawrencium	

actinoids